

Time: 3 hours

Full Marks: 300

The figures in the right-hand margin indicate marks.

Candidates should attempt Q. No. 1 from
Section – A and Q. No. 5 from Section – B
which are compulsory and three of
the remaining questions, selecting
at least one from each Section.

SECTION-A Techofworld.In

Attempt any three of the following sub-parts:

 $20 \times 3 = 60$

(a) Let $\{X_n\}$ be a sequence of random variables. If $X_n^P \to X$, where X is a random variable, then show that $g'(X_n)^P \to g(X)$ where g is a continuous function.

BD - 68/5

(Turn over)

- (b) Find the probability density function of a distribution function of a random variable whose characteristic function is defined by e-|x|. Techofworld.In
- (c) State and prove Gauss-Markov Theorem and explain its applications in linear estimation.
- (d) What is the relation between Hotelling's T² statistic and Mahalanobis D² statistic. Also, show that Hotelling T² statistic is invariant under any linear transformation.
- 2. (a) A problem in statistics is given to 3 students A, B and C whose chances of solving it are \(\frac{1}{2}\), \(\frac{3}{4}\) and \(\frac{1}{4}\) respectively. What is the probability that the problem will be solved if all of them try independently?
 - (b) A random variable X has the p. d. f. given by f(x) = 6x(1 x), $0 \le x \le 1$. Find the mean, mode and standard deviation.
 - (c) State and prove Kolmogorov's Strong Law of Large Numbers. 20×3 = 60

- 3. (a) Show that the normal distribution is a limiting form of binomial distribution.
 - (b) If X and Y are independent Gamma variates with parameters μ and ν respectively, then show that $\frac{X}{X+Y}$ is $a\beta_1(\mu,\nu)$ variate.
 - (c) The equations of two regression lines obtained in a correlation analysis are as follows:

$$3X + 12Y = 19$$
, $3Y + 9X = 46$

Obtain: Techofworld.In

- (i) The value of correlation coefficient.
- (ii) The mean values of X and Y. $20 \times 3 = 60$
- 4. (a) In a study of a random sample of 120 students, the following results are obtained:

$$\overline{X}_1 = 68$$
 $\overline{X}_2 = 70$ $\overline{X}_3 = 74$
 $s_1^2 = 100$ $s_2^2 = 25$ $s_3^2 = 81$
 $r_{12} = 0.60$ $r_{13} = 0.70$ $r_{23} = 0.65$

$$BD-68/5$$
 (3) (Turn over)

 $[s_i^2 = v(X_i)]$, where $X_{1'}$ X_2 and X_3 denote percentage of marks obtained by a student in I test, II test and the final examination respectively.

- (i) Obtain the least square regression equation of X_3 on X_1 and X_2 .
- (ii) Estimate the percentage marks of a student in the final examination, if he gets 60% and 67% in tests I and II respectively. Techofworld. In
- (b) Let X ~ N(μ, Σ), then derive the distribution of the quadratic form X' A X where A is a positive definite symmetric matrix.
- (c) Explain linear discriminant analysis and its applications in social sciences. 20×3 = 60

SECTION - B

- 5. Answer any **five** sub-questions: 12×5 = 60
 - (a) State the Cramer-Rao lower bound for the variance of an unbiased estimator. Compute

BD - 68/5

(4)

Contd.

the bound for estimation of $g(\lambda) = P[X = 0]$, where X has Poisson (λ) distribution. Does there exist an MVB estimator of $g(\lambda)$? If not, state how you would determine the UMVUE.

- (b) Define a Uniformly Most Powerful test. Derive it for testing H_0 : $\sigma^2 \le \sigma_0^2$ against H_1 : $\sigma^2 \ge \sigma_0^2$ for the variance of a normal distribution with mean 0. Derive an expression for the power function.
- (c) Stating the hypothesis, derive the stopping bounds on the sample sum for an SPRT(B, A) applied to the parameter p of a binomial distribution.
- (d) Compute the Kolmogorov-Smirnov statistic for testing the hypothesis that the following sample has come from a distribution with pdf f(x) = 2x, 0 < x < 1.
 0.22, 0.11, 0.84, 0.64, 0.42
- (e) Find an expression for the efficiency of LSD over RBD with rows as blocks.

- systematic sampling. Show that for estimating the mean, when there is a linear trend in the population, stratified sampling is n times more precise than systematic sampling.
- 6. (a) Construct the level- α UMPU test of $H_0: \sigma^2 = \sigma_0^2$ against $H_1: \sigma^2 \neq \sigma_0^2$ given a random sample the N(0, σ^2) model. Obtain the cut-off value and an expression for the power function.
 - (b) Consider the SPRT for testing the mean of $N(\mu, \sigma_0^2)$, σ_0^2 known. Derive stopping bounds on the sample sum at the n^{th} stage. Also, obtain an expression for $h(\mu) \neq 0$ satisfying $[\{f_1(x)/f_0(x)\}^{h(\mu)} f_{\mu}(x) dx = 1. \text{ Explain how you would construct the OC and ASN curves.}$
 - (c) Describe the Likelihood Ratio test procedure and state its asymptotic properties.

25+25+10 = 60

- 7. (a) Explain the regression method of estimating the population mean Y when the population mean X of the auxiliary variable is unknown.

 Derive the variance of the estimator and conditions under which it is smaller than the estimator under SRSWOR ignoring the auxiliary variable.

 Techofworld.
 - (b) Under the PPS scheme, derive a sufficient condition for WOR estimator of the population total Y_{HT} to be more efficient than the WR estimator. What is your conclusion for the equal probability sampling?
 - (c) Describe Warner's randomised response model giving an instance of its application.
 - 25+25+10 = 60
 - 8. (a) Describe a one-way random effects model and give a method of estimating the components of variance under the model.

- (b) Explain the need of confounding in factorial experiments. Setup the ANOVA table for testing the main effects of a 2⁴ factorial experiment carried out in a single replicate of 4 incomplete blocks confounding the interactions ABC and BCD.
- (c) Describe a BIBD and give an outline of its intra-block analysis. 20+20+20 = 60